JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

GGSIPU mathmatics 2011

1. Let f(x) = ax+b, a<0, then $f^{-1}(x) = f(x)$, ? x if and only if

(a) a = -1, b Č R (b) a = -1, b = 4

(c) a = -3, b Č R (d) None of these

2. The domain of $\cos^{-1}\frac{x-3}{2} - \log_{10}(4-x)$ is

(a) (1,4) (b) [1,4)

(c) (1,4] (d) [1,4]

3. If f(x) is a polynomial function of the second degree such that f(-3) = 6, f(0) = 6 and f(2) = 11, then the graph of the function f(x) cuts the ordinate x =1 at the point

(a) (1,8) (b) (1,4)

(c) (1,-2) (d) None of these

4. Let A and B be two sets, then (A Ć B)' ą (A' ą B) is equal to

(a) A' (b) A

(c) B' (d) None of these

5. The mean of 10 observations is 16.3. By an error one observation id is registered as 32 instead of 23. Then ,the correct mean is

(a) 15.6 (b) 15.4

(c) 15.7 (d) 15.8

6. Mean deviaof 6,8,12,15,10,9 through mean is

(a) 10 (b) 2.33

(c) 2 (d) None of these

7. The image of the point (2,1) w.r.t. the line x+1 = 0 is

(a) (2,5) (b) (0,5)

(c) (-4,1) (d) (-2,-3)

8. The value of x which satisfies $8^{1} + \cos x + \cos^{2} x + \ldots = 64$ in[-®,®] is

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

(a) $\pm\frac{\pi}{2}, \pm\frac{\pi}{3}$ (b) $\pm\frac{\pi}{3}$

(b) $\pm\frac{\pi}{2}, \pm\frac{\pi}{6}$ (d) $\pm\frac{\pi}{6}, \pm\frac{\pi}{3}$

9. If d = [a](a x b) +«(b x c) + ¬(cxa) is equal to and [a bc] = 1/8, then [a] + « + ¬

(a) d . (a+b+c) (b) 2d . (a+b+c)

(c) 4d . (a+b+c) (d) 8d . (a+b+c)

10. The area of the triangle formed by the points whose position vectors are 3i + j, 5i + 2j + k, i-2j+3k is

(a) $\sqrt{23}$ sq units

(b) $\sqrt{21}$ sq units

(c) $\sqrt{29}$ sq units

(d) $\sqrt{33}$ sq units

11. If (1,-2,-2) and (0,2,1) are direction ratios of two lines, then the direction cosines of a perpendicular to both the lines are

(a) $\left(\frac{1}{3}, -\frac{1}{3}, \frac{2}{3}\right)$

(b) $\left(\frac{2}{3}, -\frac{1}{3}, \frac{2}{3}\right)$

(c) $\left(-\frac{2}{3}, -\frac{1}{3}, \frac{2}{3}\right)$

(d) $\left(\frac{2}{\sqrt{14}}, -\frac{1}{\sqrt{14}}, \frac{3}{\sqrt{14}}\right)$

12. The length of the normal to the curve y = a cosh$\left(\frac{x}{a}\right)$ at any point varies as

(a) ordinate

(b) abscissa

(c) square of the abscissa

(d) square of the ordinate

13. The slope of the tangent to the curve y = $\int_0^x \frac{dx}{1+x^3}$ at the point where x =1 is

(a) ¼ (b) ½

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

(c) 1 (d) None of these

14. If f(x) = $a\log_e {}^0x^0 + bx^2+x$ has extremum at x =1 and x =3, then

(a) a -3/4, b = -1/8

(b) a = ¾, b = -1/8

(c) a = -3/4,b = 1/8

(d) None of the above

15. In the expansion of $\left(x^3 - \frac{1}{x^2}\right)^n$, n Č N, if the sum of the coefficient of x^5 and x^{10} is 0, then n is

(a) 25 (b) 20

(c) 15 (d) None of these

16. Let z_1,z_2 be two roots of the equation $z^2+az+b = 0$, z being complex number. Further assume that the origin, z_1 and z_2 form an equiliateral triangle. Then,

(a) $a^2 = b$ (b) $a^2 = 2b$

(c) $a^2 = 3b$ (d) $a^2 = 4b$

17. A square is inscribed in the circle $x^2+y^2-2x+4y-3 = 0$ with its sides parallel to the coordinate axes. One vertex of square is

(a) (3,4) (b) (3,-4)

(c) (8,-5) (d) (-8,5)

18. If f:R →R is continuous such that f(x+y) = f(x) + f(y), ? x,y ČR and f(1) = 2,then f(100) equals to

(a) 100 (b) 50

(c) 200 (d) 0

19. f(x) = x $\sin\frac{1}{x}$ is

(a) continuous but not differentiable at x= 0

(b) discontinuous but differentiable at x=0

(c) differentiable at x =0

(d) can not be determined

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

20. $\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2-bc & b^2-ca & c^2-ab \end{vmatrix}$ equals to

(a) 0

(b) 1

(c) abc

(d) (a-b) (b-c) (c-a)

21. The sum $\cos1^0 + \cos2^0 + \cos3^0 + \ldots + \cos179^0 + \cos180^0$ is equal to

(a) 0 (b) 1

(c) -1 (d) 2

22. If a,b,c are in GP and $a^{\frac{1}{x}} = b^{\frac{1}{y}} = c^{\frac{1}{z}}$, then x,y,z are in

(a) AP (b) GP

(c) HP (d) None of these

23. If A is a square marix such that $A^2 = I$, then A^{-1} is equal to

(a) I (b) O

(c) A (d) I+A

24. 5^{th} term from the end in the expansion of $\left(\frac{x^2}{2} - \frac{2}{x^2}\right)^{12}$ is

(a) $-7920x^{-4}$ (b) $7920x^4$

(c) $7920x^4$ (d) $-7920x^4$

25. Which of the following is not a logical statement?

(a) 8 is less than 6

(b) every set is a finite set

(c) Kashmir is far from here

(d) the sun is a star

26. $\tan^{-1}1 + \tan^{-2}2 + \tan^{-2}3$ is equal to

(a) 0 (b) ®

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

(c) $\frac{\pi}{2}$ (d) None of these

27. $\int_0^{\infty} \frac{1}{1+e^x} dx$ is equal to

(a) 0 (b) ®

(c) log 2 -1 (d) - log 2

28. If $^0a^0 = 8, ^0b^0 = 3$ and $^0axb^0 = 12$, then the value of a.b is

(a) 6 or -6 (b) $12\sqrt{3}$ or $-12\sqrt{3}$

(c) 8 or -8 (d) None of these

29. The value of $^nC_0 - ^nC_1 + ^nC_2 - \ldots + (-1)^n {}^nC_n$ is

(a) 1 (b) 0

(c) 2^n (d) n

30. Coefficient of variation of two distribution are 50% and 60% and their arithmetic means means are arithmetic means are 30 and 25 respectively. Difference of their standard deviation is

(a) 1 (b) 1.5

(c) 2.5 (d) 0

31. If I,j,k are the usual three perpendicular are the usual three perpendicular unit vectors then the value of i.(jxk) + j.(ixk) + k.(ixj) is

(a) 0 (b) -1

(c) 3 (d) 1

32. The solution of y dx – x dy + log x dx = 0 is

(a) y – log x -1 =Cx

(b) x + log y + 1=Cx

(c) y + log x + 1 =Cx

(d) y + log x – 1 =Cx

33. Which of the following differential equation has $y = c_1e^x + c_2e^{-x}$ as the general solution ?

(a) $\frac{d^2y}{dx^2} + y = 0$ (b) $\frac{d^2y}{dx^2} - y = 0$

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

(c) $\frac{d^2y}{dx^2} + 1 = 0$ (d) $\frac{d^2y}{dx^2} - 1 = 0$

34. $\int \frac{1}{\sin(x-a)\sin(x-b)} dx$ is equal to

(a) $\frac{1}{\sin(b-a)} \log\left|\frac{\sin(x+b)}{\sin(x+a)}\right| + C$

(b) $\frac{1}{\sin(b+a)} \log\left|\frac{\sin(x-b)}{\sin(x-a)}\right| - C$

(c) $\frac{1}{\sin(b+a)} \log\left|\frac{\sin(x-b)}{\sin(x-a)}\right| + C$

(d) None of the above

35. $\int \frac{dx}{x^2\sqrt{4-x^2}}$ is equal to

(a) $\frac{1}{4}\left(\frac{\sqrt{4-x^2}}{x}\right) + C$ (b) $\frac{1}{2}\left(\frac{\sqrt{4-x^2}}{x}\right) + C$

(c) $-\frac{1}{4}\left(\frac{\sqrt{4-x^2}}{x}\right) + C$ (d) $-\frac{1}{2}\left(\frac{\sqrt{4-x^2}}{x}\right) + C$

36. If $\tan^{-1} 2$, $\tan^{-1} 3$ are two angles of a triangle, then the third angle is

(a) 30^0 (b) 45^0

(c) 60^0 (d) 75^0

37. $\lim_{x\to 0}\left(\frac{16^x+9^x}{2}\right)^{1/x}$ is equal to

(a) 25/2 (b) 12 (c) 1 (d) ¼

38. Let a = min{x^2+2x+3, $x \in R$} and b = $\lim_{\theta\to 0}\frac{1-\cos\theta}{\theta^2}$. The value of $\sum_{r=0}^{n} a^r.b^{n-r}$ is

(a) $\frac{2^{n+1}-1}{3.2^n}$ (b) $\frac{2^{n+1}+1}{3.2^n}$

(c) $\frac{4^{n+1}-1}{3.2^n}$ (d) None of these

39. The matrix A = $\begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$ is a

(a) diagonal matrix

(b) symmetric matrix

(c) skew-symmetric matrix

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

(d) identity matrix

40. A teacher takes 3 children frpom her class to the zoo at a time as often as she can, but she does not taker the same three children to the zoo more than once. She finds that she goes to the zoo 84 times more that a particular child goes to the zoo. The number of children in her class is

(a) 12 (b) 10

(c) 60 (d) None of these

41. If A =(-3,4), B =(-1,-2), C = (5,6), D=(x,-4) are vertices of a quadrilateral such that aABD = 2 aACD. Then,x is equal to

(a) 6 (b) 9 (c) 69 (d) 96

42. The area of the parallelogram formed by the points (1,1,1), (-1,5,5), (2,2,5) is

(a) 81 (b) 9

(c) 36 (d) 18

43. If f(x) = $\frac{9^x}{9^x+3}$,then f$\left(\frac{1}{2012}\right)$ + f$\left(\frac{2}{2012}\right)$ + ... + $\left(\frac{2011}{2012}\right)$ is equal to

(a) 1005 (b) 1005.5

(c) 1006 (d) 1006.5

44. $\sqrt{1 - sin^2 101^0}$. sec 101^0

(a) 0 (b) 2

(c) -1 (d) 2

45. $\tan^{-1}\left(\frac{1}{1+2}\right) + \tan^{-1}\left(\frac{1}{1+(2)(3)}\right)$

$\tan^{-1}\left(\frac{1}{1+(3)(4)}\right)$+...+$\tan^{-1}\left(\frac{1}{1+n(n+1)}\right)$ = $\tan^{-1}$ —

(a) $\frac{n}{n+1}$ (b) $\frac{n+1}{n+2}$

(c) $\frac{n+2}{n+1}$ (d) $\frac{n}{n+2}$

46. If A_{3x3} and det (A) = 6, then det (2 adj (A) is equal to

(a) 48 (b) 8

(c) 288 (d) 12

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL

JbigDeaL JbigDeaL JbigDeaL JbigDeaL
Download from www.JbigDeaL.in Powered By © JbigDeaL™

47. The probability that a leap year o0nly 52 Sundays is

(a) $\frac{4}{7}$ (b) $\frac{5}{7}$

(c) $\frac{6}{7}$ (d) $\frac{1}{7}$

48. If $\int \frac{2^x}{\sqrt{1-4^x}}dx = {}^a\sin^{-1}(2^x) + C$, then ^{a} equals to

(a) log 2 (b) $\frac{1}{2}$ log 2

(c) $\frac{1}{2}$ (d) $\frac{1}{\log 2}$

49. If If S is circumcenter, G the centroid, O the orthocenter of aABC, then SA+SB+SC is equal to

(a) SG (b) OS

(c) SO (d) OG

50. The centre and redius of the sphere $r^2 - 2r(3i+4j-5k)+1= 0$ are

(a) 3i+4j-5k,1 (b) -3i-4j+5k,7

(c) -3i-4j+5k,7 (d) 3i+4j-5k,7

JbigDeaL JbigDeaL JbigDeaL JbigDeaL Download from www.JbigDeaL.in Powered By © JbigDeaL JbigDeaL JbigDeaL JbigDeaL JbigDeaL